

## Whey-cide Antimicrobial Solution

DESCRIPTION	Whey-cide is a peroxyacetic acid-based microbiocide developed for use in Federally Inspected dairy processing plants. If used as directed, it will reduce the incidence of bacteria and microorganisms that cause contamination, spoilage and decay, as well as pathogenic organisms.
DIRECTIONS	<p>Whey-cide use is acceptable for direct or indirect contact in or on whey and/or lactose intended for use as a food or beverage for human consumption. Since its use is intended as an indirect process additive, it does not need to be declared on any subsequent product labeling. In accordance with the FDA Food contact Notification #879, only DeLaval or its authorized distributors are allowed to use peroxyacetic acid products for whey and/or lactose production intended for human consumption.</p> <p>Dose: Maximum dose is 15 ppm (as peracetic acid) in whey and 30 ppm in Lactose</p> <p>Storage: Whey-cide™ containers must be stored in an upright position. The storage area should be well ventilated and shaded from sunlight as well as protected from sources of radiant heat. Contamination of the product, especially heavy metal ions and alkali, must be avoided.</p>
FEATURES	<ul style="list-style-type: none"><li>• Whey-cide is an equilibrium mixture of peracetic acid, hydrogen peroxide, acetic acid, proprietary ingredients, and water</li></ul>
BENEFITS	<ul style="list-style-type: none"><li>• Whey-cide exhibits excellent bactericidal and fungicidal activity against a wide range of microorganisms</li><li>• It degrades to oxygen, water, and carbon with a 1/2 life of approximately 20 minutes in whey</li></ul>
SAFETY INFORMATION	UTILIZE APPROPRIATE PERSONAL PROTECTIVE EQUIPMENT AS PER SDS. HANDLE AND STORE AS INDICATED ON PRODUCT LABEL.

IF ON SKIN: In case of contact, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Get immediate medical attention. Wash clothes before reusing.

IF IN EYES: In case of contact, immediately flush with plenty of water for at least 15 minutes. Get immediate medical attention.

IF SWALLOWED: Drink large quantities of water. Do not induce vomiting. Never give anything by mouth to an unconscious person. Get immediate medical attention.

IF INHALED: Immediately move to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult, give oxygen. Get immediate medical attention.

Consult your Safety Data Sheet for additional safety information.